Carto


CLASSIFICATION AND CORRELATION

OF

THE SOILS OF

WARREN COUNTY INDIANA

APRIL 1987



U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
MIDWEST NATIONAL TECHNICAL CENTER
LINCOLN, NEBRASKA

UNITED STATES DEPARTMENT OF AGRICULTURE Soil Conservation Service Midwest National Technical Center Lincoln, Nebraska 68508-3866

Classification and Correlation of the Soils of Warren County, Indiana

The field correlation and final field review for the soil survey of Warren County, Indiana, was held at Williamsport, Indiana, May 27-29, 1986. Participating in the final field review were James R. Barnes, soil survey party leader and William D. Hosteter, Indianapolis state office. The data reviewed consisted of the first draft of the soil survey manuscript, correlation samples, field sheets, map unit notes, laboratory data, and SCS-SOI-5 forms. Roger L. Haberman, soil correlator, MNTC, participated in the comprehensive field review on March 17-21, 1986.

The field correlation was reviewed by Roger L. Haberman in October 1986. The final correlation was base on the draft manuscript, field notes, copies of the field sheets, laboratory data, SCS-SS-6 file, selected correlation samples, and the field correlation.

Headnote for the Detailed Soil Survey Legend:

Map symbols consist of a combination of letters or of letters and numbers. The first capital letter is the initial one of the map unit name. The lowercase letter that follows separates map units having names that begin with the same letter, except that it does not separate sloping or eroded phases. The second capital letter indicates the class of slope. Symbols without a slope letter are for nearly level soils or miscellaneous areas. A final number of 2 indicates that the soil is moderately eroded and a number 3 indicates that the soil is severely eroded.

SOIL CORRELATION OF WARREN COUNTY, INDIANA

. १ . म्हिन वे राज्या राज्या स्थाप के प्रतिकार राज्या ता र स्थाप स्थाप स्थाप	CONTRACTOR OF THE CONTRACTOR O	de tare (> 1454 monteur, cam o 	e est alve sa ent mail fond automobile à fond il loir derélèpes - sière qualitératé des les sois B
symbols	unit name		Approved map unit name
	Alford silt loam, 2 to 6 percent slopes, eroded		
	Armiesburg Variant silty clay loam, frequently flooded	1	Armiesburg Variant silty clay loam, frequently flooded
	Barce silt loam, O to 2 percent slopes		
MxB2	Barce-Montmorenci silt loams, 2 to 6 percent slopes eroded	!	Barce-Montmorenci silt loams, 2 to 5 percent slopes, eroded
	Beaucoup silty clay loam, frequently flooded, undrained	•	Beaucoup silty clay loam, frequently flooded, undrained
	Beckville loam, occasionally flooded		
	Eillett sandy loam, 1 to 4 percent slopes, eroded	•	•
JaC2	Billett sandy loam, 5 to 12 percent slopes, eroded	•	to 12 percent
	Blount silt loam, 0 to 2 percent slopes		Blount silt loam, 0 to 2 percent slopes
FsC3, FtC3, FtD3, OcC2,	complex, 8 to 20	8 0 0	Boyer-Mudlavia complex, 8 to 20 percent slopes, eroded
	Prenton silt loam, 5 to 2 percent slopes		Brenton silt loam, C to 2 percent slopes

WARREN COUNTY, INDIANA -- Continued

SEN - 1888 NAME WAYS TRAIN BLOCK BOOK SHOW CHIEF BANK OF THE ETIES BEING SHOW IN	The state that the state of the	Similar and the second of the second	en Ent (Te s). Legge in men 1952 Lines wer jant, et fin Law Legen ett fin Line (19
Field symbols		cation	Approved map unit name
ReA, ReB2	Brenton silt loam, till substratum, O to 2 percent slopes	•	Brenton silt loam, till substratum, 0 to 2 percent slopes
Lm, Lo	Brenton silt loam, moderately fine substratum, 0 to 2 percent slopes		Brenton silt loam, moderately fine substratum, 0 to 2 percent slopes
MyB2, RgB2,	Xenia Variant silt Loam, 1 to 6 percent slopes, eroded	!	moderately wet, 1 to
CbA, CdA	Camden silt Loam, G to 2 percent slopes		Camden silt loam, 0 to 2 percent slopes
CbB2, CeB2	Camden silt loam, 2 to 6 percent slopes, eroded	•	*
	Camden silt loam, till substratum, 2 to 6 percent slopes, eroded	•	till substratum, 2
EuA	Carmi Loam, 0 to 2 percent slopes		Carmi loam, 0 to 2 percent slopes
Са	Comfrey loam, rarely flooded	1	Comfrey loam, stratified substratum, rarely flooded
Cs, Cu, Ot	Comfrey loam, frequently flooded	:	Comfrey loam, stratified substratum, frequently flooded, undrained
CtB2	Corwin stilt loam, 2 to 6 percent slopes, eroded		Corwin silt loam, 2 to 6 percent slopes, eroded
Cz, Rm, Ts, Wx	Cyclone silty clay Loam	Cz	Cyclone silty clay Loam

WARREN COUNTY, INDIANA -- Continued

Field symbols	unit name		Approved map unit name
	Drummer silty clay Loams		Drummer silty clay Loams
	Drummer silty clay Loam, gravelly substratum		Drummer silty clay loam, gravelly substratum
	Du Page Loam, I frequently flooded	•	Du Page Loam, frequently flooded
	Eldean gravelly loam, 2 to 6 percent slopes, eroded	:	Eldean gravelly loam, 2 to 6 percent slopes, eroded
	Eldean silt loam, 0 to 2 percent slopes	_	Eldean silt loam, G to 2 percent slopes
	Elston sandy loam, 0 to 3 percent slopes		Elston sandy loam, 0 to 3 percent slopes
	Gilboa silt loam, O to 2 percent slopes	_	Gilboa silt loam, 8 to 2 percent slopes
FoB2, FoA	:Glenhall silt loam, : sandy substratum, 1 : to 4 percent slopes, : eroded	:	: to 4 percent slopes,
FpB2, FpA, TfA, TfB	: !Glenhall silt loam, ! till substratum, 1 ! to 4 percent slopes, ! eroded	:	l till substratum, 1
₩nF	Gosport Variant shaly silt loam, 25 to 40 percent slopes	1	
HeG, HeF	Hennepin complex, 30 to 70 percent slopes		•
WnP. WnC. WnC2. OeB2	<pre>High Gap silt loam, 2 to 9 percent slopes, stony</pre>		High Gap silt loam, 2 to 9 percent slopes, stony

WARREN COUNTY, INDIANA -- Continued

Field symbols	unit name	Publi- cation symbol	unit name
	High Gap Variant Loam, 2 to 6 percent slopes, eroded		High Gap Variant Loam, 2 to 6 percent slopes, eroded
	High Gap Variant Loam, 6 to 12 percent slopes, eroded	•	High Gap Variant loam, 6 to 12 percent slopes, eroded
	Houghton muck, drained		Houghton muck, drained
	Houghton muck, undrained		Houghton muck, undrained
	Iona silt loam, 1 to 4 percent slopes, eroded	:	: 4 percent slopes,
IpA, MuS2	Ipava silt loam, n to 2 percent slopes		Ipava silt loam, ^ to 2 percent slopes
	Jules silt loam, frequently flooded		Jules silt Loam, frequently flooded
	Brenton Variant silt Loam	:	Lafayette silt loam, 2 to 2 percent slopes
La• De• Ro	La Hogue silt loam, 0 to 2 percent slopes		La Hogue silt loam, (to 2 percent slopes
·	La Hogue silt loam, till substratum, 0 to 2 percent slopes	:	La Hogue silt loam, I till substratum, O I to 2 percent slopes
	Landes-Landes Variant complex, frequently flooded	1	Landes-Chatterton complex, frequently flooded
GmB3, GLB3, MzB3	Markham silty clay Loam, 2 to 6 percent slopes, severely eroded		Markham silty clay loam, 2 to 6 percent slopes, severely eroded

WARREN COUNTY, INDIANA -- Continued

a. But the trial Mill Both on somethic this war are this pair t	Both that the part of the same are also proposed in	La Part de l'autorité des des constitues de la constitue de la	新食し (1888) - C. Good (1998) (1997) - Good (1889) - John (1997) (1888) (1997) (1998)
symbols	Field map unit name	cation	unit name
PwB2, SyB2, VaB2	Markham-Symerton silt loams, 2 to 6 percent slopes, eroded	8 6	loams, 2 to 6
	Martinsville Loam, C to 2 percent slopes		-
	Martinsville loam, 2 to 6 percent slopes, eroded	:	
	Martinsville loam, 6 to 12 percent slopes, eroded	:	Martinsville loam, 6 to 12 percent slopes, eroded
	Miami loam, 15 to 25 percent slopes, eroded	1	
MoC3	Miami clay loam. 6 to 12 percent slopes. 1 severely eroded	1	l 12 percent slopes,
stD3	Miami clay loam, 12 to 18 percent slopes, severely eroded	:	Miami clay loam, 12 to 18 percent slopes, severely eroded
Mk	: Milford silty clay : loam, pothole		<pre>!Milford silty clay ! loam, pothole !</pre>
MrA	<pre>!Millbrook silt loam, ! till substratum, 0 ! to 2 percent slopes</pre>	1	-
CtC2, MxC2,	Montmorenci-Barce complex, 6 to 12 percent slopes, eroded	:	Montmorenci-Barce complex, 6 to 12 percent slopes, eroded
•	Morley silt loam, moderately wet, 15 to 25 percent slopes, eroded	:	Morley silt loam, moderately wet, 15 to 25 percent slopes eroded

WARREN COUNTY, INDIANA -- Continued

THE LANGE WAS A RESIDENCE OF THE ROP ALL IN C. HAS MADE.	n in the second of the second	om rust of the delignation of a mark	য় কান্য প্ৰসৰ্থ হ'ব বিশ্বস্থান কৰে কৰে ভাৰত ভাৰত কৰে। তিনি কৰি স্থা নিৰ্ভিত কৰি স্থা নিৰ্ভিত কৰে প্ৰস্তাহ কৰে। আৰু
Field symbols	unit name	Publi- cation symbol	
	Morley silty clay loam, moderately wet, 6 to 12 percent slopes, severely eroded	6 6	Morley silty clay loam, moderately wet, 6 to 12 percent slopes, severely eroded
	Morley-Cadiz silt loams, moderately wet, 6 to 12 percent slopes, eroded		Morley-Cadiz silt loams, moderately wet, 6 to 12 percent slopes, eroded
	Mudlavia gravelly silt loam, 0 to 2 percent slopes, stony	1	Mudlavia gravelly silt loam. D to 2 percent slopes. stony
	Mudlavia cobbly silt loam, 2 to 4 percent slopes, eroded, stony	! !	loam, 2 to 4 percent
	substratum, 2 to 6	!	Ockley loam, sandy substratum, 2 to 6 percent slopes, eroded
	lockley silt loam, 0 to 2 percent slopes		Ockley silt loam, 0 to 2 percent slopes
	Ockley silt loam, 2 to 6 percent slopes, eroded	:	Ockley silt loam, 2 to 6 percent slopes, eroded
	Ormas loamy sand, 1 to 4 percent slopes		Ormas loamy sand, 1 to 4 percent slopes
	Oshtemo coarse sandy Loam, O to 2 percent slopes	:	Oshtemo coarse sandy Loam, O to 2 percent slopes
OsB, AgB, BuB2	<pre>!Oshtemo coarse sandy ! loam, 2 to 6 percent ! slopes</pre> !		Oshtemo coarse sandy Loam, 2 to 6 percent slopes

WARREN COUNTY, INDIANA -- Continued

	unit name		Approved map unit name
_	Peotone silty clay loam, pothole		Peotone silty clay Loam, pothole
, .	Piankeshaw Variant gravelly silt loam, rarely flooded		Piankeshaw Variant gravelly silt loam, rarely flooded
Pt	Pits, gravel	Pp	Pits, gravel
	Proctor silt loam, 0 to 2 percent slopes		Proctor silt loam, 0 to 2 percent slopes
	Proctor silt loam, 2 to 5 percent slopes, eroded	:	
	Proctor silt loam, till substratum, 0 to 2 percent slopes	:	Proctor silt loam, till substratum, 0 to 2 percent slopes
WvB2	Proctor silt loam, till substratum, 2 to 6 percent slopes, eroded	:	Proctor silt loam, till substratum, 2 to 6 percent slopes, world
Rb	Ragsdale silt loam	Rb	Ragsdale silt loam
KdA, MfA, MoA	Rainsville silt loam, 1 0 to 2 percent 1 slopes		Rainsville silt loam, 1 0 to 2 percent 1 slopes
•	Rainsville- Williamstown-Rush Variant silt loams, 2 to 6 percent slopes, eroded	:	Rainsville- Williamstown- Rockfield silt Loams, 2 to 6 percent slopes, eroded
RLA	Reesville silt loam, 6 to 2 percent slopes		Reesville silt loam, 1 0 to 2 percent 1 slopes
RvA, RuA, WuA, XeA	Rush Variant silt Loam, D to 2 percent slopes		Rockfield silt loam, U to 2 percent I slopes

WARREN COUNTY, INDIANA -- Continued

A T MIN WAS AND POST MIN. SEE THE STREET, THE WAY SEE THE PROPERTY WAS AND THE	or Tue Side in Michigan (St. 1984) Michigan (1944) Ann. Michigan (1944) Side Side (1944) Ann. Michigan (1944) Ann.		CENTED STREET BY BEEN SERVICE TO SERVICE AND THE SERVICE STREET STREET
Field symbols	unit name		Approved map unit name
RuB2, XeB2	Rush Variant silt loam, 2 to 6 percent slopes, eroded	1	Rockfield silt loam, 2 to 6 percent slopes, eroded
	Rodman gravelly loam, 25 to 60 percent slopes	•	Rodman gravelly loam. 25 to 60 percent slopes
	Rush silt loam. O to 2 percent slopes		Rush silt loam, 0 to 2 percent slopes
	Rush silt loam, 2 to 6 percent slopes, eroded	!	
Sb	Sable silty clay loam	Sb	Sable silty clay loam
	Shadeland Variant silt loam, 0 to 2 percent slopes	•	Shadeland Variant silt loam, 0 to 2 percent slopes
FdA, FdB2,	till substratum, 0 to 2 percent slopes	.	Starks silt loam, till substratum, 0 to 2 percent slopes
Ta, St	Stonelick-Moundhaven complex, frequently flooded	:	
SsB2, MpB3	Strawn clay loam, 2 to 6 percent slopes, severely eroded	:	
		:	Strawn clay loam, 6 to 12 percent slopes, severely eroded
	Symerton-Varna silt loams, 1 to 3 percent slopes	SyB	Symerton-Varna silt Loams, 1 to 3 percent slopes

WARREN COUNTY, INDIANA -- Continued

to the the person may be attempted and our major.	ALL PROPERTY IN THE PROPERTY OF THE BOX AND A STREET OF THE PROPERTY OF THE PR	MI 412-242-978-778	to their death out made made was welle price that we come made their cost, microtical artificial costs of the cost
Field symbols	unit name	Publi- cation symbol	
	Tuscola loam, till substratum, 6 to 12 percent slopes eroded		Tuscola loam, till substratum, 6 to 12 percent slopes, eroded
	Tuscola silt loam, till substratum, 2 to 6 percent slopes, eroded		Tuscola silt loam, till substratum, 2 to 6 percent slopes, eroded
0 r	Udorthents, loamy	Ud	Udorthents, loamy
	Udorthents, loamy, reclaimed	Ur	Udorthents, loamy, reclaimed
	Wakeland Variant silt Loam, occasionally flooded		Wakeland Variant silt loam, occasionally flooded
	Wallkill Variant silty clay loam		Wallkill Variant silty clay loam
	Warners Variant silty clay loam, undrained		Warners Variant silty L clay, drained
Wf	Washtenaw silt loam	Wh	Washtenaw silt loam
TeB2. WkB2.	Waupecan silt loam, moderately wet, 0 to 2 percent slopes	1	Waupecan silt loam, moderately wet, 0 to 2 percent slopes
WnE, WpD	Weikert Variant fine sandy loam, 35 to 80 percent slopes, very bouldery	1	Weitert Variant fine sandy loam, 35 to 80 percent slopes, very bouldery
BhA, EsA, EsB, EsB2,	Andres Variant- Elliott silt loams, O to 2 percent slopes	•	Williamsport-Elliott silt loams, 0 to 2 percent slopes

WARREN COUNTY, INDIANA -- Continued

Field symbols	Field map unit name	Publi- cation: symbol:	Approved map unit name
KdC2, MoC2, RgC2, RuC2, RuC3, RxC3	Williamstown- Rainsville silt Loams, 6 to 12 percent slopes, eroded		Villiamstown- Rainsville silt loams, 6 to 12 percent slopes eroded

Series Established by This Correlation:

Chatterton (type location in Warren County, Indiana)
Lafayette (type location in Warren County, Indiana)
Mudlavia (type location in Warren County, Indiana)
Rainsville (type location in Warren County, Indiana)
Rockfield (type location in Warren County, Indiana)
Williamsport (type location in Warren County, Indiana)

Series Dropped or Made Inactive:

None

Certification Statement:

The state soil scientist certifies that:

- 1. Detailed soil maps within this county have been joined.
- 2. The soil mapping of this county was completed as of December 1985.
- 3. The general soil map and detailed maps have been joined with Benton County (correlated in 1985) on the north; Tippecanoe County (soil survey in progress) on the east, Vermillion County (correlated in 1976) on the south. A soil survey is in progress in Vermillion County, Illinois, on the west. Join statements are on record.
- 4. The location of the typical pedon descriptions in this county are in soil areas using that reference name.
- 5. The interpretaions for all series used in this soil survey have been coordinated.

Verification of Exact Cooperator Names:

The following will be on the front of the publication:

United States Department of Agriculture Soil Conservation Service In cooperation with Purdue University Agricultural Experiment Station and Indiana Department of Natural Resources State Soil Conservation Board and Division of Soil Conservation

The citation in the box on the inside of the front cover will read:
"This survey was made cooperatively by the Soil Conservation Service,
Purdue University Agricultural Experiment Station, and the Indiana
Department of Natural Resources, State Soil Conservation Board, and
Division of Soil Conservation. It is part of the technical assistance
furnished to the Warren County Soil and Water Conservation District.
Financial assistance was made available by the Warren County Board of
County Commissioners."

Disposition of Original Atlas Field Sheets:

The original atlas field sheets for Warren County will be retained by the Indiana state office and will be used in the map compilation and finishing procedures. Copies have been made for fire protection purposes. The state office at Indianapolis will prepare the atlas sheets for publication by December 1986.

Prior Soil Survey Publications:

The first soil survey of Warren County was made in 1916 (ref. citation). This survey updates the first survey and provides additional information and larger maps that show the soil in greater detail.

Soil Survey of Warren County, Indiana; E. J. Grimes, Indiana Department of Geology; and E. H. Stevens, U. S. Department of Agriculture. 39 pp., illus., 1916.

Instructions for Map Finishing:

The conventional and special symbols used in this survey are listed on the attached SCS-37A. These are the only symbols that will be shown on the published maps. The maps will be finished using the "Guide for Soil Map Finishing", July 1976. Overlays and special instructions are prepared to show the separation of symbols on the field sheets.

Warren County Final Correlation

Instructions for Map Compilation and Map Finishing:

Special instructions for Map Finishing. The correlation legend will be followed except as noted on the following sheets.

Sheets 3, 10, 11, 12, 17, 19, 20, 24, 26, 41, 45, 48, 49, 53,	
43, 27, 46, 50, 18	All the Dw goes to Cz
4	All the Cz goes to Dw
11	All the Du in the area outlined goes to Cz
17	All the Sd in the area outlined goes to Cz
19	All the Sz in the area outlined goes to Cz
10, 11, 17,	All the Co, Du and Sz in the area <u>outlined</u> goes to Cz
18	All the Co, Du, and Dv goes to Cz
5, 12, 14, 19, 20	All the FdA, SlA and SnA in the area <u>outlined</u> goes to MtA
7	All the FdA, SIA and SnA outside the blue area goes to MtA
13	All the FdA, S1A and SnA on the sheet goes to MtA
5, 13	All the FpA on the sheet goes
5	All the GgA in the area outlined goes to BsA
5, 6, 13	All the FoB2 on the sheet goes to PuB2
6, 13	All the SyB2 on the sheet goes to BdB2
11, 17, 18, 24, 25, 28, 31, 32, 33, 38, 42, 54, 56	All the FsA on the sheet goes to Oca

11, 17, 18, 24, 25, 28, 31, 32, 33, 38, 42, 54, 56	All the FsB2 on the sheet goes to OcB2
5 2	All the FsB2 in the area <u>outlined</u> goes to OcB2
12, 13, 19, 20	All the GgA on the sheet goes to BsA
13, 19, 20	All the AkA, AkB, AkB2, GgB, La and Lk goes to BsA
11, 24, 25, 28 (inset), 32, 33, 38, 43, 46, 49, 52, 55 (inset), 55	All the HhB2 on the sheet goes to HfB
28, 34, 39, 54 (inset)	All the HhB2 in the area <u>outlined</u> goes to HfB
28, 33	All the WnB in the area outlined goes to HhB2
28, 39, 54 (inset)	All the HhC2 in the area outlined goes to HfB
52, 55, 55 (inset)	All the HhC2 on the sheet goes to HfB
32, 38, 52, 55 (inset), 55	All the HhA on the sheet goes to HfB
13	All the BbA, FoA and SyA goes to PuA
13, 20	All the FpB2 on the sheet goes to PuB2
21, 27, 28, 34, 39, 54 (inset)	All the WnF in the area outlined goes to WpG
38, 43, 46, 47,	All the WnF on the sheet goes to WpG
28, 34	All the WpG in the area outlined goes to GoF
33	All WpF in the area <u>outlined</u> in the lower left corner goes to WpG
3 3	All WpF on the rest of the sheet goes to GoF
25	All the RIA on the sheet

goes to S1A

41 (section 11), 42, 46, 52	All the Ao goes to Cz
45, 48, 49	All the Ao in the area outlined goes to Cz
41, 45, 48, 49, 52	All the KdB2 on the sheet goes to CaB2
4 2	All the KdB2 in the area outlined goes to CaB2
46 (section 13)	All the RuB2 goes to CaB2
42	All the KdC2 in the area outlined goes to MxC2
41	All the KdC2 on the sheet goes to MxC2
46	All the CdB2 on the sheet goes to CbB2
46 (section 24)	All the MfA goes to MdA
6	All the AkA, AkB, AkB2, GgA, GgB, La and Lk on the sheet goes to BsA (except as noted)
6	All the BbA, FoA, FpA, and SyA on the sheet goes to PuA (except as noted)
6	All the FpB2 on the sheet goes to PuB2
5	All the BbA in the area outlined goes to PuA
6, 13	All the FoB on the sheet goes to PuB2
11	All the Dv in the area <u>outlined</u> goes in Cz
1 4	All the BbA on the sheet goes to PuA
28 (Inset)	All the WpF on the inset goes to WpG
5	All the Cz in the area <u>outlined</u> goes to Dw

State:

U.S. DEPARTMENT OF AGRICULTURE SOIL CONSERVATION SERVICE

CONVENTIONAL AND SPECIAL Soil Survey Area: WARREN COUNTY SYMBOLS LEGEND

INDIANA

Date: 11/86

DESCRIPTION	SYMBOL	DESCRIPTION	SYMBOL	DESCRIPTION	SYMBOL
CULTURAL FEATUR	ES	CULTURAL FEATUR	ES (cont.)	SPECIAL SYMBOLS F SOIL SURVEY	OR
BOUNDARIES		MISCELLANGOUS CULTURAL FEATUR	ES	SOIL DELINEATIONS AND SOIL SYMBO	LS
		Farmstead, house (amit in urban areas	•	ESCARPMENTS (CeA FoB2
County or parish		Church	å	Bedrack (points down slape)	***************************************
Miner civil division		School	8	Other than bedrack (points down slope	•
				SHORT STEEP SLOPE	••••••
Field short matchine & realling					
AD HOC SOUNDARY (lase) Small airport, airfield, park, elifield complery, or fleed pool				MISCEL LANEOUS	
STATE COORDINATE TICK		WATER FEATURES			
LAND DIVISION CORNERS		DRAINAGE		Gravelly appl	••
(sections and land grants)		Perennial, dauble line			
Divided (median chown if scale per		Perennial, single line			
	•	Internution	·		
County, farm or ranch		Drainage and			
		Canals or ditches			
ROAD EMBLEMS & DESIGNATIONS					
Federal	287	Drainage and/or irrigation	•		
State	(52)			Sleny spot, very steny apot	o 33
		LAKES. PONDS AND RESERVOIRS		RECOMMENDED AD HOC SOIL SYMBOL!	5
RAILROAD	<u> </u>	Perunnus	water w		
		MISCELLANGOUS WATER FEATURES	,	Very poorly drained soils in potholes	d o
		Marsh or swamp	**		
DAMS		Wel spel	*	Very severely eroded spot	#
Stedium er small				Overwash	×
PITS					
Gravet put	×				
Mine or querry	*				

PRIME FARMLAND

(Only the soils considered prime farmland are listed. Urban or built-up areas of the soils listed are not considered prime farmland. If a soil is prime farmland only under certain conditions, the conditions are specified in parentheses after the soil name)

```
Map
                              Soil name
symboli
AfB2 : Alford silt loam, 2 to 6 percent slopes, eroded
      larmiesburg Variant silty clay loam, frequently flooded
Am
      : (where protected from flooding or not frequently flooded
      I during the growing season)
     Barce silt loam, 0 to 2 percent slopes
BBA
BdB2 | Barce-Montmorenci silt loams, 2 to 6 percent slopes,
      : eroded
      Beckville loam, occasionally flooded
3 k
     IBillett sandy loam, 1 to 4 percent slopes, eroded
3 m B 2
     iBlount silt loam, f to 2 percent slopes (where drained)
3 o A
     iBrenton silt loam, 0 to 2 percent slopes (where drained)
3rA
      IBrenton silt loam, till substratum, 0 to 2 percent slopes
3 s A
      ! (where drained)
      !Brenton silt loam, moderately fine substratum, 0 to 2
3 WA
      I percent slopes (where drained)
     ICadiz silt loam, moderately wet, 1 to 6 percent slopes,
CaB2
      : eroded
      !Camden silt loam, 0 to 2 percent slopes
CbA
      !Camden silt loam, 2 to 6 percent slopes, eroded
CP35
      1Camden silt loam, till substratum, 2 to 6 percent slopes,
CdB2
      eroded
CfA
      :Carmi loam, O to 2 percent slopes
C g
      !Comfrey loam, stratified substratum, rarely flooded
      1 (where drained)
CtB2 : Corwin silt loam, 2 to 6 percent slopes, eroded
Cz
      1Cyclone silty clay loam (where drained)
      IDrummer silty clay loams (where drained)
WC
      IDrummer silty clay loam, gravelly substratum (where
Эx
      1 drained)
Эy
      IDu Page loam, frequently flooded (where protected from
      : flooding or not frequently flooded during the growing
      : season)
Ed32
      IEldean gravelly loam, 2 to 6 percent slopes, eroded
      IEldean silt loam, 0 to 2 percent slopes
EgA
EVA
      lElston sandy loam, 0 to 3 percent slopes
      !Gilboa silt loam, 0 to 2 percent slopes (where drained)
GgA
```

SOIL SURVEY WARREN COUNTY, INDIANA

PRIME FARMLAND -- Continued

```
Soil name
Map :
sympol
3k32 :Glenhall silt loam, till substratum, 1 to 4 percent
     ! slopes, eroded
     High Gap Variant Loam, 2 to 6 percent slopes, eroded
Hh32
     !Iona silt loam, 1 to 4 percent slopes, eroded
IoB2
     lipava silt loam, 0 to 2 percent slopes (where drained)
IpA
     !Jules silt loam, frequently flooded (where protected from
Ju
     : flooding or not frequently flooded during the growing
     1 season)
     ILafayette silt loam, 3 to 2 percent slopes (where
LCA
     : drained)
Ld
     ILa Hogue silt loam. 0 to 2 percent slopes (where drained)
     La Hoque silt loam, till substratum, 9 to 2 percent
Lk
     : slopes (where drained)
     Landes-Chatterton complex, frequently flooded (where
Lp
     I protected from flooding or not frequently flooded during
     the growing season)
McB2 : Markham-Symerton silt loams, 2 to 6 percent slopes,
     ! eroded
MdA
     IMartinsville loam, 0 to 2 percent slopes
Md32 | Martinsville loam, 2 to 6 percent slopes, eroded
     iMillbrook silt loams till substratum, 0 to 2 percent
MtA
     1 slopes (where drained)
     10ckley loam, sandy substratum, 2 to 5 percent slopes,
3b32
     : eroded
     !Ockley silt loam. 0 to 2 percent slopes
O C A
Oc32 lockley silt loam, 2 to 6 percent slopes, eroded
OsA
     :Oshtemo coarse sandy loam, 0 to 2 percent slopes
     10shtemo coarse sandy loam, 2 to 6 percent slopes
3 s 3
PrA
     IProctor silt loam, 0 to 2 percent slopes
PrB2 : Proctor silt loam, 2 to 6 percent slopes, eroded
PuA : Proctor silt loam, till substratum, 0 to 2 percent slopes
    iProctor silt loam, till substratum, 2 to 6 percent
     I slopes, eroded
     !Radsdale silt loam (where drained)
₹b
     Rainsville silt loam, 6 to 2 percent slopes
RDF
    !Rainsville-Williamstown-Rockfield silt loams. 2 to 5
RfB2
      1 percent slopes, eroded
     Reesville silt loam, 3 to 2 percent slopes (where
RLA
      ! drained)
     !Rockfield silt loam, 0 to 2 percent slopes
ROA
RoB2 | Rockfield silt loam, 2 to 6 percent slopes, eroded
     Rush silt loam, 7 to 2 percent slopes
RtA
```

PRIME FARMLAND -- Continued

	中部 中部 Mail 中部 Nail 中部 Nail 中部 Nail 中部 Nail 中部 A Nail 中 A Nail h A Nai
Map	Soil name
SYMDO	L!
Rt32	Rush silt loam, 2 to 6 percent slopes, eroded
Sb	ISable silty clay loam (where drained)
SeA	IShadeland Variant silt Loam, 0 to 2 percent slopes (where
	: drained)
SLA	IStarks silt loam, till substratum, 0 to 2 percent slopes
	: (where drained)
Sr	<pre>iStonelick=Moundhaven complex, frequently flooded (where</pre>
	I protected from flooding or not frequently flooded during
	! the growing season)
Sy3	<pre>!Symerton=Varna silt loams, 1 to 3 percent slopes</pre>
Tw32	ITuscola silt loam, till substratum, 2 to 6 percent
	: slopes, eroded
₩ C	!Wakeland Variant silt loam, occasionally flooded (where
	! drained)
₩h	(Washtenaw silt Loam (where drained)
ALA	!Waupecan silt loam, moderately wet, ^ to 2 percent slopes
WrA	
	! (where drained)
- 1 170: 0:00 -0:00 1000	** OF THE PROPERTY OF THE PROP

Approved: April 29, 1987

RODNEY F. HARNER Head, Soils Staff

Midwest NTC

CONVERSION LEGEND FOR WARREN COUNTY, INDIANA

* - With white Print Print (IV) (Print (Alley & art (art	-	is the first propagation of the same con-	Control Backeton Pal Print Pal	mm sigt of garden to the gard		THE RESERVE OF THE SAME SAME SHAPE SERVE SERVE SERVE	
Field ca	bli- tion: mbol:		ubli- : ation : ymbol !	Field	Publi- : cation : symbol :		Publi- cation symbol
Am A AvA W AgB C	fB2 IrA IsB	C x A C y D w D x D b A	SLA RWA DW DX PUA	Ga Ge Gf GgB GgB2	Bk Sr Sr GgA GgA	MfB2 MfC2 MgB2 Mk MoE2	TwB2 TuC2 ObB2 Mr MoE2
AkB2 E An R AnB P	rA SA EtA	DbB2 De Df Dg Dp	PuB2 ! Ld ! Lk ! EwA ! Lk	G L B 3 G m C 3 He G Hg B 2 Hg C 2	MaB3 MwC3 HeG BdB2 MuC2	MpC3 MpD3 MrA MyC2 MyE2	MpC3 MpD3 MtA MxC2 MvE2
BbA E BpA E BrA E	Am BbA BoA BrA	Du Dv Ee EnA EnB2	DW DW BK MYA MZB2	HhB2 Ho Ho HeF HhA	HhB2 Ho Hm HeG HhB2	MzC3 McA McB3 McC2 MdC3	MwC3 SyB StB3 MxC2 MdC2
BbB2 BbC2 MBhA	dB2 1uC2	EsA EsB	CfA : EVA : WrA : WrA : WrA	IoB2	HhC2 IpA IoB2 Ju BnC2	MdD2 MaD3 MdE2 MeA MeB2	MoE2 MpD3 MoE2 McA MdB2
C a C C C C C C C C C C C C C C C C C C	a l bb A l bb B 2 l	FDB2 FsA FsB2 FsD2 FdA	GkB2 EgA EdE2 BpD2 SLA	KdB2 KdC2 KdC3	,	MfC3	R dA TuC2 GgA GgA CaB2
CtB2 CCv L	ctB2 cA cz	FoA FoB2		La Lb Lb A Lb B Lh A	Ld : Lp : Lp : Lp : Lp	MLB2 MmB2 Mn MoA MoB2	CaB2 RoB2 Bk RdA RfB2
ChC2 E CLB2 R Cn L	BpD2 RfB2 k	FpA FrA FrB2 FsC2 FsC3	SyB : McB2 : BpD2 :		Lp Lk BwA BwA	MoC2 MoC3 MoD2 MpB3 Ms	WtC2 MpC3 MoE2 StB3 SeA
CtA E CtC2 M Cu C	gA BbA MuC2 Ss SLA	FtC3 FtD3 GgA GLB2 GmB3	BpD2 BpD2 GgA CaB2 MaP3	LzB McB2 MdA MdB2 MdC2	SLA McE2 MdA MdB2 MdC2	MuB2 MxA MxB2 MxC2 MyB2	IPA BbA BdB2 MuC2 CaB2

of a limit of the last of the	WARREN COUNTY: INDIANA Continued						
	cation	Field	cation	Field	Publi- cation symbol	Field	
MyD2 MzB3 MzD3 MzE3 OcA	MaB3 MvE2 MvE2	RLA	R b R L A R p G	St StD3 SyA SyB2 SyC2	SyB McB2	WUA WUB2 WVA WVB2 WX	PuA
0 cB2 0 nB2 0 nC2 0 pB 0 r	BmB2 BnC2 OpB	RtA RtB2 RVA PVB2 ReA	RtB2 RoA RoB2			XeA XeB2	RoA RoB2
0 s B 0 u 0 c B 0 c C 2 0 d A	Ur 0 c B2 BpD2	ReB2 RgB2 RgC2 Rm Ro	WtC2 Cz	TeB2 TfA TfB ThA Ts	WLA GkB2 GkB2 OcA Cz	0 0 0 0 0 0 1 0 0	
0 dB 2 0 e A 0 eB 2 0 m A 0 mB 2	GgA HfB GgA	RsE2 RtC2 RuA RuB2 RuC2		TxA VaA VaB2 Vs Wa	PrA Sy5 McB2 Dw Wg	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
0 n A 0 n B 0 p A 0 p B 2 0 s A	0 p B	RuC3 PxC3 Sb SeA Sg	WtC2 WtC2 Sb SeA Wc	Wb Wf WnE WnF WpG	We Wh HfB GoF WpG		
OsC OsC2 Ot Pm PrA	BpD2 Cs Pm	SLA Sm Sr StB3 StC3	StA Be Sr StB3 StC3	WdA WkB2 WmA WmB2 WmC2	WLA WLA WLA BpD2		
PrB2 Pt PuA PuB2 Py	Pp PuA PuB2	SyB Sd Sh Sk Sn	SyB Dw Wc SlA SlA	WmC3 WnC WnC2 WnD WnE	BpD2 HfB HfB WpG WpG		
Pb Pg Ph Pn PwA	Hm Pm Pm Pm SyB	SnA SsB2 SsC2 SsD2 SsE2	StA StB3 StC3 MoE2 MoE2	WpD WpF Wr WsA WtA	WpG WpG Dx SLA SLA		

CLASSIFICATION OF PEDONS SAMPLED FOR LABORATORY ANALYSIS

1. Laboratory Data From NSSL With SCS-8 Forms Completed

Sampled as:	Pedon Sample Number	Publication Symbol	Approved Name
Drummer	S84IN-171-026	Dw	Drummer, till substratum
Drummer	S84IN-171-029	Dw	Drummer, till substratum
Shadeland Variant	S84IN-171-020	SeA	<pre>1/Shadeland Variant Aeric Ochraqualf; fine, mixed, mesic</pre>
High Gap Variant	S85IN-171-001	HhB2	<pre>1/High Gap Variant Ultic Hapludalf; fine-loamy, mixed, mesic</pre>
Rush Variant	S85IN-171-002	RoA <u>1</u> /	<pre>2/Rockfield</pre>
Andres Variant	S85IN-171-003	WrA <u>1</u> /	2/Williamsport

2. Laboratory Data From Purdue University with SCS-8 Forms Completed

Montmorenci	S83IN-171-1	BdB2	Montmorenci
Beecher	S83In-171-2	WrA	Beecher taxadjunct Udollic Ochraqualf; fine, mixed, mesic; inclusion in mapping unit
LaHogue	S83IN-171-3	BsA	Brenton, till substratum
Markham	S83IN-171-4	McB2	<pre>1/Markham; less clay in the C horizon than recognized</pre>
Ashkum	S83IN-171-5	Dw	Drummer, till substratum
Rush	S84IN-171-1	RtA	<pre>1/Rush; 2Bt less acid than recognized, 3 value in 3Bt outside series range</pre>

 $[\]underline{1}/$ Representative pedon for the series in Warren County

 $[\]frac{\overline{2}}{}$ Type location for series

2. <u>Laboratory Data From Purdue University (cont.</u>)

Sampled as:	Pedon Sample Number	Publication Symbol	Approved Name
Elston	S84IN-171-2	EvA	<pre>1/Elston; taxadjunct; Mollic Hapludalf</pre>
Elston	S84IN-171-3	CfA	<u>1</u> /Carmi
Fox	S84IN-171-4	EgA	Eldean taxadjunct; Typic Hapludalf; clayey-skeletal; inclusion in mapping unit
Ipava	S84IN-171-5	IpA	<u>1</u> /Ipava
Jules	S84IN-171-6	Ju	<pre>1/Jules taxadjunct; Typic Udifluvent; fine-silty, mixed, (calcareous), mesic</pre>
Oshtemo	S84IN-171-7	OsB	<pre>1/Oshtemo; Coarser sand than recognized in the series</pre>
Stonelick	S84IN-171-8	Sr	<u>1</u> /Stonelick
Ragsdale	S84IN-171-9	Rb	<pre>1/Ragsdale taxadjunct; Typic Haplaquoll; fine-silty, mixed, mesic</pre>
Sable	S84IN-171-10	Sb	Sable
Reesville	S84IN-171-11	RIA	<pre>1/Reesville; Upper Bt more acid and BC more alkaline than recognized; chroma at 6 in C and value of 6 in Bt outside series</pre>

 $[\]underline{1}$ / Representative pedon for the series in Warren County

2. <u>Laboratory Data From Purdue University (cont.</u>)

Sampled as:	Pedon Sample Number	Publication Symbol	Approved Name
Iona	S84IN-171-12	IoB2	<pre>1/Iona taxadjunct; Aquic Hapudalf; fine-silty, mixed, mesic</pre>
Alford	S84IN-171-13	AfB2	<pre>1/Alford; More alkaline in 7-11 inch layer and lower solum and 2C than recognized</pre>
Ross	S84IN-171-15	Dy	<u>1</u> /DuPage
Gessie	S84IN-171-16	Sr	Stonelick
Armiesburg	S84IN-171-17	Am	<u>1</u> /Armiesburg Variant
Hennepin	S84IN-171-18	HeG	<u>1</u> /Hennepin
Weikert Variant	S84IN-171-21	WpG	Weikert Variant; Lithic Eutrochrept; loamy-skeletal, mixed, mesic; inclusion in mapping unit
Xenia Variant	S84IN-171-22	CaB2	Cadiz taxadjunct; Aquic Hapudalf; fine-silty, mixed, mesic; inclusion in map unit
Glynwood	S84IN171-23	CaB2	Cadiz taxadjunct; Aquic Hapludalf; fine-silty, mixed, mesic; inclusion in map unit

 $[\]underline{1}/$ Representative pedon for the series in Warren County

2. Laboratory Data From Purdue University (cont.)

Sampled as:	Pedon Sample Number	Publication Symbol	Approved Name
Blount	S84IN-71-24	BoA	<pre>1/Blount; Lower solum and 2C contain less clay than recognized; lower solum more alkaline than recognized</pre>
Miami	S84IN-171-25	HeG	Miami; Less clay in the argillic horizon than recognized; inclusion in map unit
Drummer	S84IN-171-27	Dw	Drummer
Drummer	S84IN-171-28	Dw	<u>1</u> /Drummer
Eldean Variant	S84IN-171-42	MzB2	<u>1/2</u> /Mudlavia
Eldean Variant	S84IN-171-43	MyA	Mudlavia taxadjunct; Typic Hapludalf; loamy-skeletal, mixed, mesic; inclusion in map unit
Kendallville Variant	S84IN-171-44	RfB2	<u>1/2</u> /Rainsville

 $[\]underline{1}/$ Representative pedon for the series in Warren County $\underline{2}/$ Type location for series

Notes to Accompany
the Classification and Correlation
of Warren County, Indiana
by
William D. Hosteter
and Roger L. Haberman

The till sheet known in Indiana as the Snider till and which is the till of the Normal ground moraine and the Bloomington recessional moraine has a clay content which ranges from about 24 to 30 percent. Consequently, many of the pedons in Warren County in those series which are commonly associated with till having more than 27 percent clay have slightly less clay than the series range in the lower B and the C horizon.

ALFORD SERIES

These soils are more alkaline in the lower part of the solum and 2C horizon than recognized. They also have free carbonates above a depth of 80 inches. The 7-11 inch layer is more alkaline than the series range. This is likely due to liming. The thickness of the Peorian silts is less than 60 inches. This soil is mapped within 2-3 miles of the Wabash River.

ARMIESBURG VARIANT

These soils are formed in silty alluvium. About 1800 acres were mapped in Warren County.

BLOUNT SERIES

The lower 2B and 2C horizon contains less clay than the series range. The lower 2B also is less acid than the series range.

CADIZ SERIES

Value of 5 and chroma of 6 in the Bt and 2Bt; value of 5 in the 2BC; and value of 6 in the 2C are outside the series range. There is less gravel in the till than recognized. The 2C horizon contains less clay than recognized.

CAMDEN SERIES

The Bt horizon is slightly thicker than recognized for the series.

CARMI SERIES

The Bt horizon above a depth of 20 inches contains more than 15 percent gravel which is outside the series range. The C horizon typically is more alkaline than recognized.

CHATTERTON SERIES

This series is established by this correlation. The soils formed in sandy alluvium. About 600 acres were mapped in Warren County. Soils mapped as Landes Variant in Montgomery County are within the range of this series. There were 1512 acres of Landes Variant mapped in Montgomery County.

COMFREY SERIES

The Cg unit has chroma of 2 in the A horizon and value of 6 in the C horizon which are outside the series range. The Cs unit has hue of 10YR in the C horizon, which is outside the series range.

CYCLONE SERIES

A small portion (less than 200 acres) of this unit is mapped in the "heavy" till area in Warren County. In these areas the lower part of the solum and the underlying till contain more clay than the series range.

DU PAGE SERIES

The lower part of the B horizon and also the C horizon is silt loam which is outside the series range.

ELDEAN SERIES

The series is assigned very rapid permeability in the underlying material in Warren County. The series only allows rapid permeability, however, the SIR shows greater than 6 inches/hour. The coarse fragments in this soil are not dominantly limestone.

ELSTON SERIES

These soils are taxadjuncts to the series because base saturation is too low for a Mollisol. The soils are Mollic Hapludalfs. A footnote should be added to the tables stating that this soil is a source of soil and gravel at depths greater than 80 inches.

GLENHALL SERIES

The type location of this series is moved to Warren County. The concept is changed from the substratum being sand and gravel to stratified silty and loamy material with thin sandy strata.

GOSPORT SERIES

Soils called Gosport Variant in the field correlation are correlated as Gosport following review of base saturation on associated soils.

These soils have slightly more coarse fragments than recognized for the series.

HIGH GAP SERIES

The 2Bt is silty clay loam which is outside the series range. It contains more than 15 percent fine sand or coarser.

HIGH GAP VARIANT

These soils formed in outwash and residuum from shale. Involved is about 670 acres.

IONA SERIES

These soils are taxadjuncts to the series because mottles are within the upper 10 inches of the argillic horizon. They are Aquic Hapludalfs.

IPAVA SERIES

The solum is slightly thinner than the series range. Depth to free carbonates is also less.

JULES SERIES

These soils are taxadjuncts to the series because they contain more clay in the control section. They are fine-silty.

LAFAYETTE SERIES

This series is established by this correlation. The soils formed in silty material and the underlying loamy and gravelly outwash. About 930 acres were mapped in Warren County. About 800 acres of this soil was correlated as Brenton Variant in Montgomery County. This soil is also being mapped in Tippecanoe County.

MARKHAM SERIES

Permeability will be narrowed to moderately slow in the C horizon so that it is compatible with other soils formed in similar material. This soil contains less clay in the C horizon than the series range. Also 3 chroma in the A horizon of the severely eroded unit is outside the series range.

MONTMORENCI SERIES

These soils are taxadjuncts because they do not have chroma 2 mottles within the upper 10 inches of the argillic horizon. They are Mollic Hapludalfs.

MORLEY SERIES

The C horizon typically contains less clay than the series range. Value of 5 and chroma of 3 in the A horizon are outside the series range.

MUDLAVIA SERIES

This series is established by this correlation. The soils formed in gravelly loamy outwash. About 2,700 acres were mapped in Warren County.

OCKLEY SERIES

Value of 5 and chroma over 3 in the 3Bt are outside the series range.

ORMAS SERIES

The soils lack a Bt horizon and have a slightly thicker 2Bt than recognized.

OSHTEMO SERIES

The sand size fraction in the soils is coarse. Except for the A horizon, this is outside the series range.

PIANKESHAW VARIANT

These soils formed in gravelly and stony colluvium and are calcareous to the surface. Involved is 200 acres.

RAGSDALE SERIES

These soils are taxadjuncts to the series because they do not have sufficient clay increase for an argillic horizon. The soils are Typic Haplaquolls.

RAINSVILLE SERIES

This series is established by this correlation. The soils formed in a thin layer of silty material, loamy outwash and glacial till. About 11,000 acres were mapped in Warren County.

REESVILLE SERIES

The upper part of the Bt is more acid than recognized. The Bt horizon has a value of 6 which is outside the series range. The BC horizon is more alkaline than recognized and chroma of 6 in the C horizon is outside the series range.

ROCKFIELD SERIES

This series is established by this correlation. The soils formed in silty material, loamy outwash and till. About 11,000 acres were mapped in Warren County.

RUSH SERIES

The 2Bt and 3Bt horizons are less acid than recognized. Value of 3 in the 3Bt horizon is outside the series range.

SABLE SERIES

Lab data for the typical pedon shows 4.6 percent CaCO, in the 30-45 inch layer but no effervescence was noted in the field.

SHADELAND VARIANT

These soils formed in outwash and dominantly residuum from shale. Involved is abut 900 acres.

STRAWN SERIES

The lower part of the Bt horizon is loam which is outside the series range.

TUSCOLA SERIES

These soils are taxadjuncts to the series because gray mottles are not within the upper 10 inches of the argillic horizon. They are Typic Hapludalfs.

VARNA SERIES

The C horizon contains slightly less clay than the series range.

WAKELAND VARIANT

These soils formed in silty and loamy alluvium. They have free carbonates at a depth of 20 to 40 inches. Involved is about 1,120 acres.

WALLKILL VARIANT

These soils formed in sediments washed from adjacent slopes and deposited over organic deposits in potholes. Involved is about 320 acres.

WARNERS VARIANT

These soils formed in lacustrine sediments. Involved is about 225 acres.

WASHTENAW SERIES

These soils are taxadjuncts to the series because they contains less sand. They are fine-silty.

WAUPECAN SERIES

The solum is slightly thicker and depth to carbonates is slightly greater than recognized.

WEIKERT VARIANT

These soils formed in loamy residuum from sandstone. Involved is about 860 acres.

WILLIAMSPORT SERIES

This series is established by this correlation. These soils formed in silty material, loamy outwash, and glacial till. Involved is about 8,200 acres.

WILLIAMSTOWN SERIES

These soils are taxadjuncts to the series because they do not have gray mottles within the upper 10 inches of the argillic horizon. They are Typic Hapludalfs.

CLASSIFICATION OF THE SOILS

(An asterisk in the first column indicates a taxadjunct to the series. See notes for a description of those characteristics of this taxadjunct that are outside the range of the series)

	10 10 11 12 13 13 14 15 15 15 15 15 15 15
Soil name :	Family or higher taxonomic class
:	
	Fine-silty, mixed, mesic Typic Hapludalfs
	Fine, mixed, mesic Fluventic Hapludolls
Variant :	
	Fine-loamy, mixed, mesic Typic Argiudolls
	Fine-silty, mixed, mesic Fluvaquentic
	Haplaquolls
	Coarse-loamy, mixed, nonacid, mesic Aquic
	Udifluvents
	Coarse-loamy, mixed, mesic Mollic Hapludalfs
	Fine, illitic, mesic Aeric Ochraqualfs
-	Coarse-loamy, mixed, mesic Typic Hapludalfs
	Fine-silty, mixed, mesic Aquic Argiudolls
	Fine-silty, mixed, mesic Typic Hapludalfs
	Fine-silty, mixed, mesic Typic Hapludalfs
	Coarse-loamy, mixed, mesic Typic Hapludolls
	Sandy, mixed, mesic Fluventic Hapludolls
-	Fine-loamy, mixed, mesic Cumulic Haplaquolls
	Fine-loamy, mixed, mesic Typic Argiudolls
	Fine-silty, mixed, mesic Typic Argiaquolls
	Fine-silty, mixed, mesic Typic Haplaquolls Fine-loamy, mixed, mesic Cumulic Hapludolls
	Fine, mixed, mesic Typic Hapludalfs
	Fine, illitic, mesic Aquic Argiudolls
	Coarse-loamy, mixed, mesic Typic Argiudolls
	Fine-loamy, mixed, mesic Aquic Argiudolls
	Fine-loamy, mixed, mesic Mollic Hapludalfs
	Fine, illitic, mesic Typic Dystrochrepts
· · · · · · · · · · · · · · · · · · ·	Fine-loamy, mixed, mesic Typic Eutrochrepts
	Fine-loamy, mixed, mesic Typic Hapludalfs
	Fine-loamy, mixed, mesic Ultic Hapludalfs
Variant :	tano adamy, manda, modeo de
	Euic, mesic Typic Medisabrists
	Fine-silty, mixed, mesic Typic Hapludalfs
	Fine, montmorillonitic, mesic Aquic Argiudolls
	Coarse-silty, mixed (calcareous), mesic Typic
	Udifluvents
	Fine-loamy, mixed, mesic Aquic Argiudolls
,	The second of th

CLASSIFICATION OF THE SOILS -- Continued

```
Soil name : Family or higher taxonomic class
THE REST OF THE RE
 Lafayette---: Fine-silty, mixed, mesic Aquic Argiudolls
 Landes----: Coarse-loamy, mixed, mesic Fluventic Hapludolls
 Markham----! Fine, illitic, mesic Mollic Hapludalfs
 Martinsville : Fine-loamy, mixed, mesic Typic Hapludalfs
 Miami----: Fine-loamy, mixed, mesic Typic Hapludalfs
 Milford----! Fine, mixed, mesic Typic Haplaquolls
 Millbrook----! Fine-silty, mixed, mesic Udollic Ochraqualfs
*Montmorenci : Fine-loamy, mixed, mesic Aquollic Hapludalfs
 Morley----: Fine, illitic, mesic Typic Hapludalfs
 Moundhaven---: Sandy, mixed, mesic Typic Udifluvents
 Mudlavia----: Clayey-skeletal, mixed, mesic Typic Hapludalfs
 Ockley----: Fine-loamy, mixed, mesic Typic Hapludalfs
 Ormas----! Loamy, mixed, mesic Arenic Hapludalfs
 Oshtemo----: Coarse-loamy, mixed, mesic Typic Hapludalfs
 Peotone----! Fine, montmorillonitic, mesic Cumulic
                          : Haplaquolls
 Piankeshaw | Loamy-skeletal, mixed (calcareous), mesic Typic
     Variant ! Udifluvents
 Proctor----! Fine-silty, mixed, mesic Typic Argiudolls
*Ragsdale----: Fine-silty, mixed, mesic Typic Argiaquolls
 Rainsville---: Fine-loamy, mixed, mesic Typic Hapludalfs
 Reesville----! Fine-silty, mixed, mesic Aeric Ochraqualfs
 Rockfield----: Fine-silty, mixed, mesic Typic Hapludalfs
 Rodman----: Sandy-skeletal, mixed, mesic Typic Hapludolls
 Rush----: Fine-silty, mixed, mesic Typic Hapludalfs
 Sable----: Fine-silty, mixed, mesic Typic Haplaquolls
 Shadeland : Fine, mixed, mesic Aeric Ochraqualfs
     Variant
 Starks----: Fine-silty, mixed, mesic Aeric Ochraqualfs
 Stonelick----: Coarse-loamy, mixed (calcareous), mesic Typic
                        ! Udifluvents
  Strawn----: Fine-loamy, mixed, mesic Typic Hapludalfs
  Symerton----: Fine-loamy, mixed, mesic Typic Argiudolls
*Tuscola----: Fine-loamy, mixed, mesic Aquic Hapludalfs
                           ! Loamy, Udorthents
  Udorthents
  Varna----: Fine, illitic, mesic Typic Argiudolls
                           1 Coarse-silty, mixed, nonacid, mesic Aeric
  Wakeland
      Variant
                                Fluvaquents
  Wallkill
                           Fine, mixed, mesic Cumulic Haplaquolis
      Variant
  Warners
                           : Fine-silty, mixed, mesic Fluvaquentic
      Variant
                                Haplaquolls
```

SOIL SURVEY WARREN COUNTY, INDIANA

CLASSIFICATION OF THE SOILS -- Continued

. A PAGE TAND THREE TIME STATE STATE STATE STATE STATE STATE AND THREE TAND STATE ST	数 1990 数元子 高級 1980 1980 1980 1980 1980 1980 1980 1980
Soil name	Family or higher taxonomic class
1	
}	
	Fine-loamy, mixed, nonacid, mesic Aeric Fluvaquents
Waupecan	Fine-silty, mixed, mesic Typic Argiudolls
•	Coarse-loamy, mixed, mesic Dystric Eutrochrepts
Williamsport	Fine, mixed, mesic Aquic Argiudolls
· ·	Fine-Loamy, mixed, mesic Aquic Hapludalfs
THE MEDITION OF THE PERSON OF	TO THE STATE OF TH